arXiv:1604.06688v2 [math.GT] 8 Jul 2016

INTERSECTION NORMS ON SURFACES AND BIRKHOFF CROSS SECTIONS

MARCOS COSSARINI AND PIERRE DEHORNOY

ABSTRACT. For every finite collection of curves on a surface, we define an associated (semi-)norm on the first homology group of the surface. The unit ball of the dual norm is the convex hull of its integer points. We give an interpretation of these points in terms of certain coorientations of the original collection of curves. Our main result is a classification statement: when the surface has constant curvature and the curves are geodesics, integer points in the interior of the dual unit ball classify isotopy classes of Birkhoff cross sections for the geodesic flow (on the unit tangent bundle to the surface) whose boundary is the symmetric lift of the collection of geodesics. Birkhoff cross sections in particular yield open-book decompositions of the unit tangent bundle.

CONTENTS

Introduction 1
1. Intersection norms 7
2. Unit balls and coorientations 9
2.a. Coorientations and signed intersections 10
2.b. Eulerian coorientations 10
2.c. Eikonal functions 11
2.d. Proof of Theorem B 13
3. Birkhoff cross sections with antithetic boundary for the geodesic flow 13
3.a. Constructions of Birkhoff cross sections with antithetic boundary 14
3.b. Asymptotic cycles and existence of sections 15
3.c. Classes of surfaces with given boundary 17
3.d. Proofs of Theorems C and D 18
4. Questions 20
References 22

INTRODUCTION

This article has two goals. First we introduce an elementary family of norms, called *intersection norms*, on the first homology group of a real surface and we study their unit balls. These norms can be seen as (parametrized) 2-dimensional analogs of the Thurston norm on the second homology group of a 3-manifold. Secondly, we use these intersection norms to classify up to isotopy certain 2-dimensional objects in some 3-manifolds, namely, Birkhoff cross sections with

Date: June 12, 2016.

prescribed boundary for the geodesic flow in the unit tangent bundle to a negatively curved surface (or to the torus with a flat metric).

Intersection norms. Let Σ be a real compact surface with empty boundary. A **wall system** on Σ is a finite, self-transverse system of unoriented compact curves immersed in Σ, each of them an immersed circle. The wall system can be seen as a graph embedded in Σ, with vertices of even degree. The geometric intersection number of two multi-curves is usually defined as the minimum of the number of intersection points of two representants of the free homotopy classes of the two multi-curves that have disjoint double points. We propose here a variation of this notion where we fix one multi-curve (the wall system), but minimize over the homology class of the second.

So let γ denote a fixed wall system on Σ. For α a path on Σ, the geometric intersection $i_\gamma(\alpha)$ is the minimal number of intersection points with γ of a multi-curve homotopic to α and in general position with respect to γ. Beware that this definition is not symmetric since the wall system γ is fixed and not allowed to change in its homotopy class. Given a homology class a in $\mathrm{H}_1(\Sigma;\mathbb{Z})$, we then minimise the intersection number over all closed multi-curves in a. This defines a function $\|\cdot\|_\gamma : \mathrm{H}_1(\Sigma;\mathbb{Z}) \to \mathbb{N}$ by

$$\|a\|_\gamma := \min_{[\alpha]=a} i_\gamma(\alpha) = \min_{\substack{[\alpha]=a \\ \alpha \pitchfork \gamma}} |\{\alpha \cap \gamma\}|.$$

Theorem A. *Let Σ be a compact oriented surface and γ a wall system on Σ. The function $\|\cdot\|_\gamma$ extends uniquely to a continuous function $\|\cdot\|_\gamma : \mathrm{H}_1(\Sigma;\mathbb{R}) \to \mathbb{R}_+$, which is convex, and linear on rays from the origin. If, furthermore, the wall system γ fills Σ in the sense that the complement $\Sigma \setminus \gamma$ is the union of topological discs, then $\|\cdot\|_\gamma$ is a norm.*

Theorem A is an exact transposition of W. Thurston's result defining a norm on the second homology group of a 3-manifold [Thu86, Thm 1]. In particular the (semi-)norm $\|\cdot\|_\gamma$ has the property of taking integer values on integer classes. Thurston showed [Thu86, Thm2] that this implies that the unit ball, denoted here by $B_{\|\cdot\|_\gamma}$, is very peculiar: it is a polyhedron with finitely many sides, which are all given by linear equations with integer coefficients. Equivalently, this means that the closed unit ball of the dual norm on $\mathrm{H}^1(\Sigma;\mathbb{R})$, denoted here by $B^*_{\|\cdot\|_\gamma}$, is the convex hull of finitely many integer points.[1]

In the case of the Thurston norm, some of these extremal points of the dual ball could be interpreted using Euler classes of fibrations on the circle [Thu86, Thm 3]. An interpretation of all extremal points was then given in terms of Euler class of taut foliations by D. Gabai (unpublished, see [Yaz16, Thm 3.3]), and in terms of flows by D. Calegari [Cal06].

The analog statement for intersection norms is simpler. Considering a wall system γ as a graph whose vertices are the double-points and whose edges are the simple arcs of γ, a **coorientation** of γ is the a choice of a coorientation for every edge of γ. A given wall system has only finitely many coorientations. A coorientation is **Eulerian** if around every double point, there are two positively and two negatively cooriented edges. A coorientation ν can be paired with an oriented curve α using signed intersection. If ν is Eulerian, it turns out that the pairing $\nu(\alpha)$ depends only on the homology class of α, so that an Eulerian coorientation ν induces an integral cohomology class $[\nu] \in$

[1] Thurston's proof of this fact is not as natural as one might expect. M. De La Salle recently gave a more direct proof [Sal16].

$\mathrm{H}^1(\Sigma;\mathbb{Z})$.[2] One can wonder which classes are represented by such Eulerian coorientations. A first remark is that, representing a class a by a curve α which minimises the geometric intersection with γ, one sees that $|\nu(a)|$ is not larger than $\|a\|_\gamma$. A second remark is that the parity of $\nu(a)$ is fixed by γ: indeed, since all intersection points are counted with a coefficient ± 1, the parity of $\nu(\alpha)$ is determined by the parity of $i_\gamma(\alpha)$ and does not depend on ν; since γ is a graph of even degree, the parity of $i_\gamma(\alpha)$ does not change if we replace α by a homologous curve. Our second result states that these restrictions are the only ones: the classes of the Eulerian coorientations are exactly the integer points in $B^*_{\|\cdot\|_\gamma}$ that are *congruent to* $[\gamma]_2$ *mod 2*. More interestingly, the extremal points of $B^*_{\|\cdot\|_\gamma}$ correspond to some Eulerian coorientations.

Theorem B. *Let Σ be a compact oriented surface and γ a wall system on Σ. The dual unit ball $B^*_{\|\cdot\|_\gamma}$ in $\mathrm{H}^1(\Sigma;\mathbb{R})$ is the convex hull of the points in $\mathrm{H}^1(\Sigma;\mathbb{Z})$ given by all Eulerian coorientations of γ. Equivalently, for every a in $\mathrm{H}_1(\Sigma;\mathbb{Z})$, we have*

$$\|a\|_\gamma = \min_{[\alpha]=a} i_\gamma(\alpha) = \max_{\substack{\nu \text{ Eulerian} \\ \text{coor. of } \gamma}} \nu(a).$$

Moreover every point in $B^*_{\|\cdot\|_\gamma} \cap \mathrm{H}^1(\Sigma;\mathbb{Z})$ that is congruent to $[\gamma]_2 \bmod 2$ is the class of some Eulerian coorientation (see Figure 1).

Not only does this result provide an interpretation of the integer points inside the unit ball of the dual norm, it also gives an effective way of computing the norm $\|\cdot\|_\gamma$, since it reduces the minimisation over an infinite number of curves into a maximisation over a finite number of coorientations.

Classification of Birkhoff cross sections for geodesic flows. Let M be a compact, orientable smooth n-manifold without boundary, and let X be a non-singular vector field on M. In order to understand the dynamics of X it is desirable to find a **global cross section** for (M, X), namely a compact, orientable hypersurface S without boundary such that

- S is embedded in M,
- S is transverse to X,
- every orbit of X intersects S after a bounded time: we have $\phi^{[0,T]}(S) = M$ for some $T > 0$.

When such a section exists, there is a well-defined first-return map on S and the first-return time is bounded from above by definition and from below by compactness. In this case the manifold M fibers over the circle with fiber S. The pair (M, X) is homeomorphic to $(S \times [0,1]/_{(p,1)\sim(f(p),0)}, \tau_p \frac{d}{dz})$, where τ_p is the first-return time on S and $\frac{d}{dz}$ denotes the vector field tangent to the $[0,1]-$coordinate. The dynamics of X is then, up to the time-reparametrisation function τ, the dynamics of the first-return map f on S.

A standard argument shows that two global sections are isotopic if and only if they are homologous. Indeed the flow then realizes the isotopy between such homologous sections (see for example the discussion at the beginning of [Thu86, Section 3]). Therefore questions of existence and classification of global sections are of algebraic nature. Indeed, a necessary and sufficient

[2] According to the universal coefficients theorem for cohomology, for any abelian group G, the cohomology group $\mathrm{H}^1(\Sigma, G)$ is naturally isomorphic to the group $\mathrm{Hom}(\mathrm{H}_1(\Sigma;\mathbb{Z}); G)$ of group homomorphisms from $\mathrm{H}_1(\Sigma;\mathbb{Z})$ to G. However, in this article we can take this identification as the definition of $\mathrm{H}^1(\Sigma; G)$, since the usual definition of cohomology groups (as homology groups of the singular cochain complex) won't be used.



FIGURE 1. Illustration of Theorems B and D. On the left a wall system γ consisting of four geodesics on the torus T^2, and an Eulerian coorientation (blue arrows). On the right the dual unit ball $B^*_{\|\cdot\|_\gamma} \subset \mathrm{H}^1(\mathrm{T}^2, \mathbb{R})$ of the associated intersection norm. The empty circle denotes the origin. The big dots denote those classes in $\mathrm{H}^1(\mathrm{T}^2, \mathbb{Z})$ congruent to $[\gamma]_2$ mod 2. Among these classes, 10 (in blue, green an red) are in the dual unit ball $B^*_{\|\cdot\|_\gamma}$ and correspond to all cohomology classes of Eulerian coorientations of γ (Theorem B). For example, the class corresponding to the blue coorientation is the blue point. The blue and green points lie in the interior of $B^*_{\|\cdot\|_\gamma}$, hence describe the two isotopy classes of Birkhoff cross sections for φ_{geod} bounded by $-\overleftrightarrow{\gamma}$, while the 8 red points are on the boundary of $B^*_{\|\cdot\|_\gamma}$ and describe isotopy classes of surfaces transverse to φ_{geod}, but not intersecting every orbit, and bounded by $-\overleftrightarrow{\gamma}$ (Theorem D).

condition for a given homology class σ in $\mathrm{H}_2(M;\mathbb{Z})$ to contain a global section has been described by S. Schwartzmann and F. Fuller: the set of *Scharzmann asymptotic cycles* [Sch57] in $\mathrm{H}_1(M;\mathbb{R})$ has to lie in the half-space $\{\langle\sigma,\cdot\rangle > 0\}$, where $\langle\cdot,\cdot\rangle$ denotes the algebraic intersection pairing $\mathrm{H}_2(M;\mathbb{R}) \times \mathrm{H}_1(M;R) \to \mathbb{R}$. This implies for example that vector fields on $\mathbb{S}^3$ never admit global sections. Further results of W. Thurston [Thu86] and D. Fried [Fri82] imply that in the case of a pseudo-Anosov flow, the set of homology classes of global sections is an open cone with finitely many extremal rays.

For Σ a Riemannian surface, the unit tangent bundle $\mathrm{T}^1\Sigma$ is the subset of $\mathrm{T}\Sigma$ of norm 1-vectors. It is a 3-manifold whose points are of the form (p, v) for p a point of Σ and v a tangent vector at p of norm 1. The geodesic flow φ_{geod} on $\mathrm{T}^1\Sigma$ is the vector field whose orbits are lifts of geodesics: for g an arbitrary geodesic of Σ travelled at speed 1, the orbit of φ_{geod} going through $(g(0), \dot{g}(0))$ is given by $\varphi^t_{\mathrm{geod}}(g(0), \dot{g}(0)) = (g(t), \dot{g}(t))$. The geodesic flow on a negatively curved surface has been studied since Hadamard who remarked its sensibility to initial condition [Had1898]. It even became the paradigm of 3-dimensional chaotic systems when Anosov showed its hyperbolic character [Ano67]. In general the geodesic flow depends heavily on the metric given on the surface. However Gromov remarked [Gro76] that the geodesic flows corresponding to any two negatively curved metrics on a surface are actually *topologically conjugated*, meaning that there is a homeomorphism of the tangent bundle sending the oriented orbits the first on the oriented orbits of the second. This is a consequence of the structural stability of Anosov flows. Therefore, as long as we

are only interested in the topological properties of the orbits, one can speak of *the* geodesic flow on a negatively curved surface.

Since the antipodal map $(p, v) \mapsto (p, -v)$ preserves the geodesic flow, its set of asymptotic cycles is symmetric with respect to the origin in $H_1(T^1\Sigma; \mathbb{R})$, so that geodesic flows never admit global sections.

In order to make it useful, a relaxation of the notion of global section is desirable. For M a real compact, oriented 3-manifold and X a non-singular vector field on M, a **Birkhoff cross section** for (M, X) is compact orientable surface S *with* boundary such that

- S is embedded in M,
- the interior $int(S)$ is transverse to X,
- the boundary ∂S is tangent to X,
- every orbit of X intersects S after a bounded time: we have $\phi^{[0,T]}(S) = M$ for some $T > 0$.

The third condition implies that the boundary of S is the union of finitely many periodic orbits of X. The second and third condition may look hard to realize at the same time, but actually it is not the case: in a flow box oriented so that the vector field is vertical, the general picture of a Birkhoff cross section near its boundary is that of a helicoidal staircase.

Since the interior of a Birkhoff cross section S is transverse to X, it is cooriented by X. Since M is oriented, this induces an orientation on S, and in turn an orientation of ∂S. On the other hand, ∂S is a collection of periodic orbits of X, so it is oriented by X. For every component of ∂S, these two orientations may coincide or be opposed. We say that S is a **positive Birkhoff cross section** if they coincide for every boundary component, **negative** if they are opposed (on the left), and **mixed** if they sometime agree and sometime disagree. If the fourth condition is not satisfied, namely of some orbits do not intersect the surface, we simply speak of a **transverse surface**.



negative *positive*

It turns out that Birkhoff cross sections exist much more often than global sections. In particular H. Poincaré noticed that the geodesic flow on a sphere often admits an annulus as Birkhoff cross section. This remark was generalized by G. Birkhoff who gave a family of Birkhoff cross sections for the geodesic flow [Bir17] (popularized in [Fri83]). Birkhoff's example was then given another presentation by M. Brunella [Bru94, Description 2]. Our first result is a generalization of Birkhoff's and Brunella's examples.

For γ an unoriented collection of geodesics on a surface Σ, we denote by $\overleftrightarrow{\gamma}$ the **amphithetic lift** of γ in $T^1\Sigma$, that is, the set of unit tangent vectors based on γ and tangent to γ. The set $\overleftrightarrow{\gamma}$ forms a link that is invariant by the involution $(p, v) \mapsto (p, -v)$. It is the union of $2|\gamma|$ periodic orbits of φ_{geod}, each component being oriented by the flow.

Theorem C. *Let Σ be a compact oriented Riemannian surface and γ a finite collection of closed geodesics on Σ. There is canonical a map S^{BB} (for Birkhoff-Brunella) that associates to every Eulerian coorientation ν of γ an oriented surface $S^{BB}(\nu)$ in $T^1\Sigma$ which is positively transverse to the geodesic flow and whose oriented boundary is $-\overleftrightarrow{\gamma}$. For every ν, the Euler characteristic of $S^{BB}(\nu)$ is minus twice the number of double points of γ.*

If two Eulerian coorientations ν_1, ν_2 of γ are cohomologous, then the associated surfaces $S^{BB}(\nu_1)$ and $S^{BB}(\nu_2)$ are isotopic (fixing their common boundary).

The main interest of this new construction is that it actually gives a description of all isotopy classes of negative Birkhoff cross sections with boundary $\overleftrightarrow{\gamma}$, instead of one with the previously known constructions:

Theorem D. *Let Σ be a torus with a flat metric or a higher genus-surface with a negatively curved metric. Let γ be a finite collection of closed geodesics on Σ. Then the map $[\nu] \mapsto \{S^{BB}(\nu)\}$ is a one-to-one correspondance between integer points in the closed unit ball $B^*_{\|\cdot\|_\gamma}$ congruent to $[\gamma]_2 \bmod 2$ and isotopy classes of surfaces in $\mathrm{T}^1\Sigma$ transverse to φ_{geod} with oriented boundary $-\overleftrightarrow{\gamma}$.*

The restriction of this map to the integer points in the open unit ball $\mathrm{int}(B^*_{\|\cdot\|_\gamma})$ induces a one-to-one correspondance with isotopy classes of negative Birkhoff cross sections.

Theorem D implies that the collection $\overleftrightarrow{\gamma}$ bounds a negative Birkhoff cross section if and only if the polyhedron $B^*_{\|\cdot\|_\gamma}$ contains an integer point congruent to $[\gamma]_2 \bmod 2$ in its interior. This is the case for most choices of γ, but not for all. For example, if there is a closed curve that intersects γ once or zero times, then $\overleftrightarrow{\gamma}$ does not bound a Birkhoff cross section for φ_{geod}.

Remark 1. It may look strange to deal with negative Birkhoff cross sections and not with positive ones, *i.e.,* with surfaces such that the orientation of the boundary inherited from the orientation of the surface (itself inherited from the coorientation of the interior surface by the flow) is opposed to the direction of the flow. The reason is that there is actually no positive Birkhoff cross section for the geodesic flow. One could then look at *mixed* sections, namely transverse surfaces some of whose boundary components are positively tangent to φ_{geod} and some others are negatively transverse. It is likely that there are more mixed sections than negative. We do not have analogs of Theorems C and D in this more general case, namely we do not have any elementary way to construct them all.

Remark 2. A Birkhoff surface for the geodesic flow in $\mathrm{T}^1\Sigma$ bounded by $\overleftrightarrow{\gamma}$ is a global section for the restriction of the flow to $\mathrm{T}^1\Sigma \setminus \overleftrightarrow{\gamma}$. The assumption that the boundary is $-\overleftrightarrow{\gamma}$ can be seen as a restriction on the homology class of the section: it has to lie in a certain affine subspace of $\mathrm{H}_2(\mathrm{T}^1\Sigma, \overleftrightarrow{\gamma}; \mathbb{Z})$ (see Section 3.c). On the other hand, as explained before, the geodesic flow on $\mathrm{T}^1\Sigma$ for Σ a hyperbolic surface is of Anosov type. Its restriction to $\mathrm{T}^1\Sigma \setminus \overleftrightarrow{\gamma}$ is then of pseudo-Anosov type, with singularities along the removed orbits. Thurston fibered faces Theory [Thu86, Section 3] then says that the homology classes of global sections to such a flow (and therefore of isotopy classes) is a cone in $\mathrm{H}_2(\mathrm{T}^1\Sigma, \overleftrightarrow{\gamma}; \mathbb{R})$ whose extremal rays are directed by integral vectors. D. Fried [Fri82] gives an algorithm to explicitly compute these vectors, starting from a Markov partition of the flow. So one deduces directly that the set of negative Birkhoff cross sections is given by the intersection of a cone with an affine plane: it is a polyhedron. However, the determination of this polyhedron using Fried's approach requires an explicit Markov partition for the geodesic flow on $\mathrm{T}^1\Sigma \setminus \overleftrightarrow{\gamma}$, which does not exist yet. Theorem D can be rephrased by saying the Thurston's fibered face corresponding to the geodesic flow on $\mathrm{T}^1\Sigma \setminus \overleftrightarrow{\gamma}$ is a multiple of $B^*_{\|\cdot\|_\gamma}$. So the interest of our paper lies in the elementary and explicit characters of all constructions.

Remark 3. Another possible interest of the paper is that it suggests that there may exist an object that would describe all Birkhoff cross sections for a given flow simultaneously (this role is played here by the intersection norm $\|\cdot\|_\gamma$), in the same spirit as Ghys proved [Ghy09] that Gauss linking forms describe all linking numbers between periodic orbits (and even invariant measures) for a vector field in a homology sphere.

Acknowledgments. The author Pierre D. thanks Étienne Ghys and Adrien Boulanger for many related discussions, and Elena Kudryavtseva who initiated this article by asking several questions about Birkhoff cross sections.

1. Intersection norms

In this section we define intersection norms and prove Theorem A. All statements are transcriptions of results of Thurston [Thu86] to the 2-dimensional context of a surface with a wall system on it. Although the original 3-dimensional proofs are rather easy, their transcriptions are even more elementary.

For the whole section we fix a compact surface Σ of genus g without boundary, and a wall system γ on Σ.

Given a closed multi-curve α transverse to γ and such that the multiple points of α and γ are disjoint, there is a finite number of intersection points between α and γ. What we do here is to minimize it over the homology class of α:

Definition 4. (see Figure 2) The wall system γ being fixed on Σ, the function $\|\cdot\|_\gamma : \mathrm{H}_1(\Sigma;\mathbb{Z}) \to \mathbb{N}$ is defined by

$$\|a\|_\gamma := \min_{[\alpha]=a} i_\gamma(\alpha) = \min_{\substack{[\alpha]=a \\ \alpha \pitchfork \gamma}} |\alpha \cap \gamma|.$$

Since the number of intersection points is an integer, the lower bound is always realized and $\|\cdot\|_\gamma$ takes integral values. A multi-curve that realizes the minimum is declared $\|\cdot\|_\gamma$**-minimizing**.

The function $\|\cdot\|_\gamma$ has two properties that will turn it into a semi-norm, namely it is linear on rays and convex. To prove the first point we need an elementary remark. Let us recall that a multi-curve is **simple** if it has no double points, that is, if it is an embedding.

Lemma 5 (simplification)**.** *For every wall system γ in Σ and for every class a in $\mathrm{H}_1(\Sigma;\mathbb{Z})$, there exists a $\|\cdot\|_\gamma$-minimizing multi-curve in a that is simple.*

Proof. Starting from an arbitrary α_0 in a that is minimizing, we can smooth the double points of α_0 away from γ



thus turning α_0 into a new multi-curve α which is simple. The two multi-curves are in general not homotopic, but they are homologous, hence the result. □

Lemma 6 (linearity on rays)**.** *For every a in $\mathrm{H}_1(\Sigma;\mathbb{Z})$ and for all $n \in \mathbb{Z}$ one has*

$$\|n \cdot a\|_\gamma = |n|\,\|a\|_\gamma.$$



FIGURE 2. A genus 3 surface with a collection γ made of four closed curves (black). On the left the curve α_1 (orange and bold) is transverse to γ and intersects it three times. On the right α_2 (red) is homologous to α_1 since their difference bounds a subsurface, namely the right hemisurface. The curve α_2 intersects γ only once. This number cannot be reduced to 0 in the same homology class, hence α_2 is $\|\cdot\|_\gamma$-minimizing and we have $\|(\|_\gamma[\alpha_1]) = \|(\|_\gamma[\alpha_2]) = i_\gamma(\alpha_2) = 1$.

Proof. Since one does not change the number of intersection points by reversing the orientation of a curve, one has $\|-a\|_\gamma = \|a\|_\gamma$.

We then assume $n \geq 0$. Given $a \in \mathrm{H}_1(\Sigma;\mathbb{Z})$, consider a minimizing multi-curve α in a. Since n parallel copies of α intersect γ in $n\,\|a\|_\gamma$ points, we have $\|n \cdot a\|_\gamma \leq n\,\|a\|_\gamma$.

For the other inequality, consider a multi-curve $\alpha^{(n)}$ that minimizes $\|n\cdot a\|_\gamma$. By the simplification Lemma 5, we can suppose $\alpha^{(n)}$ simple. Since $\alpha^{(n)}$ is homologous to n copies of α, its number of crossings (counted with signs) with any generic loop is a multiple of n. So, starting from an arbitrary region in the complement $\Sigma \setminus \alpha^{(n)}$ that we color with the label 0, we can color the other regions with the labels $0, 1, \dots, n-1$ in such a way that the color increases by 1 mod n when one crosses an arc of $\alpha^{(n)}$ positively (from right to left). Therefore $\alpha^{(n)}$ is the union of the n simple multi-curves α_i, each such α_i consisting on the components of $\alpha^{(n)}$ that run leaving the regions labelled i on their right, and the regions labeled $i+1$ on their left. Since they pairwise bound a subsurface of Σ, all of these n multi-curves are homologous. These implies that $\alpha^{(n)}$ is homologous to n copies of any α_i. Since it is also homologous to n copies of α, and $H_1(\Sigma;\mathbb{Z})$ has no torsion, we conclude that each α_i is homologous to α. Then it has at least $\|a\|_\gamma$ intersecions with γ, which implies that $\alpha^{(n)}$ has at least $n\|a\|_\gamma$ intersections, concluding the proof that $\|n \cdot a\|_\gamma \geq n\,\|a\|_\gamma$. □

Lemma 7 (convexity)**.** *For every a, b in $\mathrm{H}_1(\Sigma;\mathbb{Z})$ one has*

$$\|a+b\|_\gamma \leq \|a\|_\gamma + \|b\|_\gamma.$$

Proof. The union of two multi-curves that realize $\|a\|_\gamma$ and $\|b\|_\gamma$ crosses γ in $\|a\|_\gamma + \|b\|_\gamma$ points, giving $\|a+b\|_\gamma \leq \|a\|_\gamma + \|b\|_\gamma$. □

Proof of Theorem A. Every class in $\mathrm{H}_1(\Sigma;\mathbb{Q})$ is of the form $\frac{1}{q}\,a$ with $a \in \mathrm{H}_1(\Sigma;\mathbb{Z})$ and $q \in \mathbb{N}^*$. We then define $\|\frac{1}{q}\,a\|_\gamma$ as $\frac{1}{q}\|a\|_\gamma$, and the linearity on rays (Lemma 6) ensures that this definition

does not depend on the choice of q and a and that it yields a well-defined function (also denoted by $\|\cdot\|_\gamma$) from $H_1(\Sigma;\mathbb{Q})$ to $\mathbb{Q}_+$ that is linear on rays. Now convexity (Lemma 7) implies that this function extends uniquely to a convex function from $H_1(\Sigma;\mathbb{R})$ to $\mathbb{R}_+$. Indeed the extension can be defined by taking the convex hull of the epigraph (what lies above the graph), or, more precisely, the supremum of the linear functions that are smaller than $\|\cdot\|_\gamma$. The extension (still denoted by $\|\cdot\|_\gamma$) is also convex and linear on rays, hence it is a semi-norm on $H_1(\Sigma;\mathbb{R})$.

If the collection γ decomposes Σ into simply-connected regions, then γ intersects every curve that is not null-homotopic at least once. This implies that $\|\cdot\|_\gamma$ is at least 1 on non-zero integral homology classes, hence $\|\cdot\|_\gamma$ is positive on $H_1(\Sigma;\mathbb{R}) \setminus \{0\}$. Therefore $\|\cdot\|_\gamma$ is a norm. □

Remark 8. One can easily extend the notion of intersection norms to surfaces with boundary, by allowing wall systems to contain arcs with endpoints on the boundary of the surface. One then obtains two norms on $H_1(\Sigma;\mathbb{R})$ and $H_1(\Sigma,\partial\Sigma;\mathbb{R})$, depending whether one considers absolute or relative homology classes. All statements can be translated in this context.

Remark 9. One can wonder how the intersection norms compare with other known norms on the first homology of a surface. For example the *stable norm* x_g is defined in terms of a metric g by $x_g(a) = \liminf_{n\to\infty} \min_{\alpha^{(n)}\in a^n} g(\alpha^{(n)})/n$. When g is negatively curved, the stabilisation is not necessary, so that $x_g(a) = \min_{\alpha\in a} g(\alpha)$. One can check that if $(\gamma_k)_{k\in\mathbb{N}}$ is a sequence of filling geodesics for g, meaning that the sequence of invariant measures on $T^1\Sigma$ that are concentrated on the lift $\vec{\gamma}_k$ tends in the weak sense to the Liouville measure defined by g on $T^1\Sigma$, then the rescaled norms $\frac{1}{g(\gamma_k)}x_{\gamma_n}$ tend to the stable norm of g. Equivalently, the rescaled unit balls $g(\gamma_k)B_{x_{\gamma_k}}$ tend to the unit ball of the stable norm.

2. Unit balls and coorientations

For the whole section we fix a surface Σ of genus at least 1 and a wall system γ on it. The norm $\|\cdot\|_\gamma$ defined in the previous section has a very peculiar property: it takes integral values on integral classes. This property is shared for example by the ℓ_1- and ℓ_∞-norms on $\mathbb{R}^d$, whose unit balls are polyhedral. Moreover all faces of these unit balls are of the form $\{(x_1,\dots,x_d) \mid \sum x_i y_i = 1\}$ for some $(y_1,\dots y_d) \in \mathbb{Z}^d$. This is not a coincidence as was remarked by Thurston.

Theorem 10 (Thm 2 of [Thu86])**.** *If N is a seminorm on $\mathbb{R}^d$ taking integral values on $\mathbb{Z}^d$, then there is a finite subset F of $\mathbb{Z}^d$ such that $N(x) = \max_{y\in F}\langle x,y\rangle$ for all x in $\mathbb{R}^d$.*

Let us recall that a norm N on a vector space induces a dual norm N^* on the dual by $N^*(y) = \max_{x\in B}\langle x,y\rangle$ where B denotes the unit ball of N. Thurston's result can be restated by saying that the unit ball of the dual norm is the convex hull of finitely many integral points.

In our context, denote by $\|\cdot\|_\gamma^*$ the norm on $H_1(\Sigma;\mathbb{R})^* \simeq H^1(\Sigma;\mathbb{R})$ dual to $\|\cdot\|_\gamma$, by $B_{\|\cdot\|_\gamma}$ the unit ball of $\|\cdot\|_\gamma$, and by $B^*_{\|\cdot\|_\gamma}$ the unit ball of $\|\cdot\|_\gamma^*$. A direct consequence of Theorem 10 is

Corollary 11 (see Figure 3)**.** *For Σ a compact surface and γ a wall system on it, the unit ball $B^*_{\|\cdot\|_\gamma}$ is the convex hull in $H^1(\Sigma;\mathbb{R})$ of finitely many points that belong to $H^1(\Sigma;\mathbb{Z})$.*

A natural question is whether the vertices of $B^*_{\|\cdot\|_\gamma}$ (or equivalently the faces of $B_{\|\cdot\|_\gamma}$) have a nice interpretation. For example in the context of the Thurston norm, the vertices correspond



FIGURE 3. A torus with a collection γ (black) made of four curves, two vertical and two horizontal. The curve α (red and bold) intersects γ in 10 points. It is the best for a curve whose homology class is $(4, 1)$. The norm $\|\cdot\|_\gamma$ is actually given by $\|(\|_\gamma p, q) = 2|p| + 2|q|$ in the canonical coordinates. The unit balls $B_{\|\cdot\|_\gamma}$ (bold) and $B^*_{\|\cdot\|_\gamma}$ (dotted) are shown on the right. The faces of $B_{\|\cdot\|_\gamma}$ are defined by integral equations while the vertices of $B^*_{\|\cdot\|_\gamma}$ belong to $\mathbb{Z}^2$, as predicted by Thurston's result.

to the Euler classes of certain taut foliations (Gabai, see [Yaz16, Thm 3.3]), or of certain vector fields [Fri79, Mos92, Cal06]. Here we also have such an interpretation in terms of Eulerian coorientations (Theorem B), and it is the goal of this section to prove it.

2.a. **Coorientations and signed intersections.** Recall that the wall system γ is assumed to be self-transversely immersed with only double points. We denote by $V(\gamma)$ the set of double points, that we call **vertices** of γ. Consequently we denote by $E(\gamma)$ the set of connected components of $\gamma \setminus V(\gamma)$, that we call **edges** of γ. This turns γ into a graph of degree 4 embedded in Σ.

Definition 12. For e an edge of γ, a **coorientation on** e is the choice of one of the two possible ways of crossing e: from left to right, or from right to left. A **coorientation on** γ is a the choice of a coorientation for every edge in $E(\gamma)$.

There are $2^{|E(\gamma)|}$ coorientations of γ. A coorientation ν may be evaluated on an oriented immersed curve α transverse to γ: one counts $+1$ for every intersection point of α with γ if the orientation of α coincides with the coorientation of the edge, and -1 if the orientations disagree. Denoting by $\nu(\alpha)$ this intersection pairing, one sees that $\nu(\alpha)$ is an integer satisfying $|\nu(\alpha)| \leq i_\gamma(\alpha)$.

2.b. **Eulerian coorientations.** The question now is whether the above inequality may be turned into an equality for $\|\cdot\|_\gamma$-minimizing curves on the one hand, and whether $\nu(\alpha)$ may depend only on the homology class of α so that one can compute ν on a single representative. Both questions admit a positive answer if we restrict to some special coorientations, called Eulerian.

Definition 13. A coorientation on γ is **Eulerian** (or **closed**) if it vanishes on boundaries, that is, if for every region $D \subset \Sigma$ whose boundary is transverse to γ one has $\nu(\partial D) = 0$. The set of all global Eulerian coorientations is denoted by $\mathcal{E}ulCo(\gamma)$.

The set $\mathcal{E}ulCo(\gamma)$ is an affine subspace of $\{-, +\}^{E(\gamma)}$. Actually the closing condition is local: for ν to be Eulerian it is enough that around every vertex of γ there are as many positively cooriented edges than negatively cooriented. Hence, up to rotation, there are only two types (locally, at each vertex) of Eulerian coorientations:



When one travels straight along γ and encounters a vertex of the first type the coorientation changes, hence the name. For the second type on the other hand, it is as if the coorientation does not see the vertex.



Example 14. If $[\gamma]_2 \in H^1(\Sigma;\mathbb{Z}/2\mathbb{Z})$ is zero (meaning that every closed curve intersects γ an even number of times), then the regions of $\Sigma \setminus \gamma$ can be colored in black and white in such a way that adjacent regions have different colors. In this case we can coorient all edges toward the white regions. The obtained global coorientation is Eulerian.



Example 15. There always exist global Eulerian coorientations, even when $[\gamma]_2 \in H^1(\Sigma;\mathbb{Z}/2\mathbb{Z})$ is not zero. Indeed if the wall system γ is the immersion of c curves, it admits at least the 2^c Eulerian coorientations obtained by choosing a coorientation for every component and having only transparent vertices.

Lemma 16. *If ν is an Eulerian coorientation of γ, then for every multi-curve α, the pairing $\nu(\alpha)$ depends only of the homology class $[\alpha] \in H_1(\Sigma;\mathbb{Z})$.*

Proof. If two multi-curves α, α' are homologous, then their difference bounds a singular subsurface in Σ. Definition 13 implies that the pairing of the boundary of the image of a surface with an Eulerian coorientation is zero. Hence $\nu(\alpha - \alpha') = 0$, so $\nu(\alpha) = \nu(\alpha')$. □

Lemma 16 states that every Eulerian coorientation ν induces a well-defined cohomology class $[\nu]$ in $H^1(\Sigma;\mathbb{Z})$. We denote by $[\mathcal{E}ulCo(\gamma)]$ the subset of $H^1(\Sigma;\mathbb{Z})$ formed by the classes of global Eulerian coorientations on γ. Note that the class of an Eulerian coorientation is easily computed since it is enough to evaluate its pairing with $2g$ curves that generate the homology of Σ. Moreover, Eulerian coorientations give lower bounds on $\|\cdot\|_\gamma$:

Lemma 17 (Eulerian orientations are in the dual ball)**.** *For every Eulerian coorientation ν of γ and for every a in $H_1(\Sigma;\mathbb{Z})$, we have $\nu(a) \leq \|a\|_\gamma$.*

Proof. Let α be a curve in a that realizes $\|a\|_\gamma$. Then $\nu(\alpha)$ counts every intersection point of α and γ with a coefficient ± 1, while $\|a\|_\gamma$ counts all these intersection points with a coefficient $+1$, hence the inequality. □

2.c. **Eikonal functions.** An Eulerian coorientation is analog to a certain 1-form on the graph dual to γ in Σ. As such it can be seen as the differential of a certain multivariate function or, equivalently, as the projection to Σ of the differential of a function defined on the universal cover of Σ. This approach is useful for constructing Eulerian coorientations with a prescribed cohomology class, as is needed for proving Theorem B.

We consider for every path α in Σ, the number $\mathrm{Len}_\gamma(\alpha)$ of intersections with γ; this notion of length determines a (not positive definite) distance d_γ on Σ. Two points x, y are **neighbors** if $d_\gamma(x, y) = 1$.

Choose a basepoint $p_0 \in \Sigma$ and construct the universal cover $\tilde{\Sigma}$ of Σ as the set of homotopy classes $x = \{\alpha\}$ of curves α that begin at p_0 and end at any point $p =: \pi\{\alpha\}$, in particular, let x_0 be the class of the constant curve.

Lift γ to a wallsystem $\tilde{\gamma}$ in $\tilde{\Sigma}$. Any closed curve β in Σ based at p_0 determines a deck transformation $T_{\{\beta\}} : x = \{\alpha\} \in \tilde{\Sigma} \mapsto x' = \{\beta \cdot \alpha\}$, where $\beta \cdot \alpha$ is the concatenation of α after β. The deck transformation $T_{\{\beta\}}$ preserves the distance $d_{\tilde{\gamma}}$.

An Eulerian coorientation ν on M gives rise to a function $f_\nu : \tilde{\Sigma} \setminus \tilde{\gamma} \to \mathbb{Z}$ by the formula $f_\nu\{\alpha\} = \int_\alpha \nu$ consisting in counting with signs the intersection points of α with γ. This function is **eikonal**, meaning that $|f_\nu(x) - f_\nu(x')| = 1$ whenever $d_\gamma(x, x') = 1$ (and also $f_\nu(x) = f_\nu(x')$ whenever $d_\gamma(x, x') = 0$). It is also $[\nu]$-equivariant, meaning that $x' = T_{\{\beta\}}x$ implies $f_\nu(x') - f_\nu(x) = [\nu]([\beta])$.

Definition 18. A function f defined on a subset D of $\tilde{\Sigma} \setminus \tilde{\gamma}$ is said **pre-eikonal** if it satisfies $|f(y') - f(y)| \leq d_{\tilde{\gamma}}(y', y)$ and $f(y') - f(y) \equiv d_{\tilde{\gamma}}(y', y) \bmod 2$ for every $y, y' \in \tilde{\Sigma}$.

Observe that a function defined on all of $\tilde{\Sigma} \setminus \tilde{\gamma}$ is eikonal if and only if it is pre-eikonal.

Lemma 19 (Extension)**.** *Every pre-eikonal function $f : D \to \mathbb{Z}$ extends to an eikonal function $\overline{f} : \tilde{\Sigma} \setminus \tilde{\gamma} \to \mathbb{Z}$.*

Proof. (Based on footnote of [Whi34].) To define $\overline{f}(x)$, we first observe that it must lie in the interval $[f(y) - d_{\tilde{\gamma}}(x, y), f(y) + d_{\tilde{\gamma}}(x, y)]$ for every $y \in D$. So we can define $\overline{f}(x)$ as the highest common point $\overline{f}(x) := \min_{y \in D} f(y) + d_{\tilde{\gamma}}(x, y)$ of these intervals, after checking that they do have a common point, because they intersect pairwise. And indeed they do, for otherwise there would exist two points y, y' in S such that $f(y) + d_{\tilde{\gamma}}(x, y) < f(y') - d_{\tilde{\gamma}}(x, y')$, which implies $f(y') - f(y) > d_{\tilde{\gamma}}(x, y) + d_{\tilde{\gamma}}(x, z) \geq d_{\tilde{\gamma}}(y, y')$, a contraction to pre-eikonality.



We claim that the extension $\overline{f}$ is pre-eikonal (and therefore eikonal, since it is defined in all of $\Sigma \setminus \gamma$).

Indeed, to prove that $|f(x') - f(x)| \leq d_{\tilde{\gamma}}(x, x')$, it is enough to check that

$$\left|(f(y) + d_{\tilde{\gamma}}(x', y)) - (f(y) + d_{\tilde{\gamma}}(x, y))\right| \leq d_{\tilde{\gamma}}(x, x')$$

for each y, which follows from the triangle inequality in the form $|d_{\tilde{\gamma}}(x', y) - d_{\tilde{\gamma}}(x, y)| \leq d_{\tilde{\gamma}}(x, x')$. To prove that $f(x') - f(x) \equiv d_{\tilde{\gamma}}(x, x')$ modulo 2, we write

$$\begin{aligned} f(x') - f(x) &= (f(y') + d_{\tilde{\gamma}}(x', y')) - (f(y) + d_{\tilde{\gamma}}(x, y)) \text{ for certain } y, y' \in D \\ &\equiv d_{\tilde{\gamma}}(y, y') + d_{\tilde{\gamma}}(x', y') - d_{\tilde{\gamma}}(x, y) \bmod 2 \text{ since } f \text{ is pre-eikonal} \\ &\equiv d_{\tilde{\gamma}}(y, y') + d_{\tilde{\gamma}}(x', y') + d_{\tilde{\gamma}}(x, y) \text{ since plus and minus coincide mod 2} \\ &\equiv d_{\tilde{\gamma}}(x, x') \text{ since homotopic paths have congruent length mod 2.} \end{aligned}$$

□

Note that a pre-eikonal function admits in general several eikonal extensions. The one we picked in the proof is the *highest* one. It has the advantage of admitting a closed definition.

2.d. **Proof of Theorem B.** By Lemma 17, for every Eulerian coorientation ν, the class $[\nu]$ belongs to $B^*_{\|\cdot\|_\gamma}$, so we have $[\mathcal{E}ulCo(\gamma)] \subset B^*_{\|\cdot\|_\gamma}$. Conversely, by Thurston's Theorem, extremal points of $B^*_{\|\cdot\|_\gamma}$ belong to $\mathrm{H}^1(\Sigma;\mathbb{Z})$. Therefore it is enough to show that for every integer point n in the closed ball $B^*_{\|\cdot\|_\gamma} \subset \mathrm{H}^1(\Sigma;\mathbb{R})$ that is congruent to $[\gamma]_2$ mod 2 there exists an Eulerian coorientation whose cohomology class is n. This is the content of Lemma 22 below.

Let D be the orbit of x_0 by the deck action. For every closed curve α based at p_0 and for $y = \{\alpha\} \in D$ we set $f_n(y) := n([\alpha])$.

Lemma 20. *The function $f_n : D \to \mathbb{Z}$ is a n-equivariant pre-eikonal function.*

Proof. Let y, y' be two points in D that we write as $y = \{\alpha\}$ and $y' = T_{\{\beta\}}(y) = \{\beta\cdot\alpha\}$ for some closed curves α, β based at p_0. By definition we have $f_n(y') - f_n(y) = n([\beta\cdot\alpha]) - n([\alpha]) = n([\beta])$, so f_n is n-equivariant. Furthermore, if we choose β in the form $\beta = \alpha\cdot\beta'\cdot\alpha^{-1}$ with β' of minimum length, that is, $\mathrm{Len}_\gamma(\beta') = d_{\tilde\gamma}(y, y')$, we see that $|f_n(y') - f_n(y)| = |n([\beta'])| \le \mathrm{Len}_{\tilde\gamma}(\beta') = d_{\tilde\gamma}(y, y')$. Finally we have $f_n(y) - f_n(y') = n([\beta']) \equiv \mathrm{Len}_\gamma(\beta') = d_{\tilde\gamma}(y', y) \bmod 2$. Therefore f_n is pre-eikonal. □

By the Extension Lemma 19, we can extend f_n to an eikonal function $\overline{f_n}$. We chose $\overline{f_n}$ as in the proof, namely by the formula $\overline{f_n}(x) = \min_{y\in D} f_n(y) + d_{\tilde\gamma}(x, y)$.

Lemma 21. *The function $\overline{f_n}$ is n-equivariant.*

Proof. If $x = \{\alpha\}$ and $x' = T_{\{\beta\}}(x) = \{\beta\cdot\alpha\}$, then to prove that $f_n(x') - f_n(x) = n([c])$ we just need to observe that the function $f_n : D \to \mathbb{Z}$, as seen from x', looks the same, but $n([\beta])$ units higher, than as seen from x. More precisely, the contribution $f(y) + d_{\tilde\gamma}(x, y)$ of each $y = \{\alpha\} \in D$ to the formula $\overline{f_n}(x) = \min_{y\in D} f(y) + d_{\tilde\gamma}(x, y)$ is $n([\beta])$ units less than the contribution $f_n(y') + d(x', y')$ of its image $y' = T_{\{\beta\}}(y)$ to the formula $\overline{f_n}(x') = \min_{y'\in D} f_n(y') + d(x', y')$, because $f_n(y') - f_n(y) = n([\beta])$ and $d_{\tilde\gamma}(x', y') = d_{\tilde\gamma}(x, y)$. □

Lemma 22. *There is a unique Eulerian coorientation ν on γ whose lift $\tilde\nu$ satisfies $\int_b \tilde\nu = \overline{f_n}(x') - \overline{f_n}(x)$ whenever $x = \{\alpha\}$ and $x' = \{\beta\cdot\alpha\}$. This ν satisfies $[\nu] = n$.*

Proof. Since $\overline{f_n}$ is an eikonal function, there exists a unique coorientation $\tilde\nu$ of $\tilde\gamma$ whose integral on each path equals the variation of $\tilde\nu$. To prove that it descends to a coorientation ν of γ, we only need to check that it is invariant by deck transformations. Indeed if x and z are neighbors, and x', z' are the respective images via a deck transformation $T_{\{\beta\}}$, then $\tilde\nu(x', z') = \overline{f_n}(z') - \overline{f_n}(x') = (\overline{f_n}(z) + n([\beta])) - (\overline{f_n}(x) + n([\beta])) = \overline{f_n}(z) - \overline{f_n}(x) = \tilde\nu(x, z)$, as required. Finally, to see that $[\nu] = n$, note that if β is a closed loop in Σ based at a point p, and α is a curve from p_0 to p, then both the startpoint x_0 and the endpoint $x = T_{\{\alpha\cdot\beta\cdot\alpha^{-1}\}}(x_0)$ of the loop $\alpha\cdot\beta\cdot\alpha^{-1}$ are in D, and we have

$$\int_\beta \nu = \int_{\alpha\cdot\beta\cdot\alpha^{-1}} \nu = \overline{f_n}(T_{\{\alpha\cdot\beta\cdot\alpha^{-1}\}}x_0) - \overline{f_n}(x_0) = n([\alpha\cdot\beta\cdot\alpha^{-1}]) = n([\beta]).$$

□

3. Birkhoff cross sections with antithetic boundary for the geodesic flow

In this part, we make an additional assumption: now Σ denotes a Riemannian surface that may be a torus with constant curvature or a higher-genus surface with strictly negative curvature. The collection γ now consists of finitely many periodic geodesics on Σ.

In this setting, the geodesic flow $(\varphi^t_{\text{geod}})_{t\in\mathbb{R}}$ on the unit tangent bundle $\mathrm{T}^1\Sigma$ is the flow whose orbits are lifts of geodesics. Namely for g a geodesic parametrized at speed one, the orbit of φ_{geod} going though the point $(g(0), \dot g(0)) \in \mathrm{T}^1\Sigma$ is $\varphi^t_{\text{geod}}((g(0), \dot g(0)) = (g(t), \dot g(t))$. For every oriented periodic geodesic g on Σ, there is one periodic orbit of φ_{geod} corresponding to the oriented lift of g and denoted by $\vec{g}$. Then if g now denotes an unoriented geodesic on Σ, there are two associated periodic orbits of φ_{geod}, one for each orientation. We denote by $\overleftrightarrow{g}$ the union of these two periodic orbits, it is an oriented link in $\mathrm{T}^1\Sigma$ that is invariant under the involution $(p, v) \mapsto (p, -v)$. A link of the form $\overleftrightarrow{g}_1 \cup \cdots \cup \overleftrightarrow{g}_k$ is called an **antithetic link**.

Let us recall from the introduction that, given a complete flow $(\phi^t)_{t\in\mathbb{R}}$, a compact surface S with boundary is **transverse** to ϕ^t if its interior is transverse to the orbits of the flow and its boundary is the union of finitely many periodic orbits[3]. A **Birkhoff cross section** for ϕ^t is then a transverse surface S that intersects every orbit of ϕ^t. A small analysis and a compactness argument show that around the boundary S necessarily looks like a helix, so that the first-return time on $\mathrm{int}(S)$ is bounded.

In this section, we give a construction that associates to every Eulerian coorientation a surface transverse to the geodesic flow (3.a). Then we recall some facts on the existence of global sections for vector fields (3.b), before making some elementary algebraic topology for describing homology classes of surfaces with boundary (3.c). Finally we put pieces together to prove that the construction actually exhaust all possible surfaces, thus proving Theorems C and D (3.d).

3.a. **Constructions of Birkhoff cross sections with antithetic boundary.** We now explain how to associate to every Eulerian coorientation of γ a surface bounded by $\overleftrightarrow{\gamma}$ and transverse to φ_{geod}, thus proving the first part of Theorem C.

From now on we fix a global coorientation ν (not yet Eulerian) of γ. For every edge e of γ (*i.e.* segment between two double points), we consider the set $R^{e,\nu}$ of those tangent vectors based on e and pairing positively with ν. This is an rectangle in $\mathrm{T}^1\Sigma$ of the form $e \times [-\pi, \pi]$ (see Figure 4). Is is bounded by the two lifts of e in $\mathrm{T}^1\Sigma$ (called the **horizontal part** of $\partial R^{e,\nu}$) and two halves of the fibers of the extremities of e (called the **vertical part** of $\partial R^{e,\nu}$). Note the interior of $R^{e,\nu}$ is transverse to the geodesic flow φ_{geod} while the horizontal part of $\partial R^{e,\nu}$ is tangent to it. We then orient $R^{e,\nu}$ so that φ_{geod} intersects it positively. One checks that then the induced orientation on $\partial R^{e,\nu}$ is opposite to the one given by φ_{geod}.

Consider now the 2-dimensional complex $S^\times(\nu)$ that is the union of the rectangles $R^{e,\nu}$ for all edges e of γ.

Lemma 23. *The 2-complex $S^\times(\nu)$ described above has boundary $-\overleftrightarrow{\gamma}$ if and only if the coorientation ν is Eulerian.*

Proof. Since $S^\times(\nu)$ is the union of one rectangle per edge of γ, the horizontal boundary of $S^\times(\nu)$ is always $\overleftrightarrow{\gamma}$. Since the orientation is opposite to the geodesic flow, it is actually $-\overleftrightarrow{\gamma}$.

What we have to check is that the vertical boundary is empty if and only if ν is Eulerian. At every double point v of γ there are four incident rectangles, corresponding to the four adjacent edges. Now the vertical boundary of a rectangle $R^{e,\nu}$ is oriented upwards (that is, trigonometrically) at the right extremity of e (when cooriented by ν) and downwards at the left extremity. Then the vertical

[3]Often in the literature a transverse surface is only defined locally. The condition we add here on the boundary is not standard. However we keep the name for avoiding a heavier expression.



FIGURE 4. Bottom: an edge e of γ and a coorientation ν on it. Top: the corresponding rectangle $R^{e,\nu}$ in $\mathrm{T}^1\Sigma$. The dotted lines represent the fibers of some points of Σ, that is, each point on these lines represent a unit tangent vector to Σ. Since the fibers are actually circles, the top and bottom extremities of the dotted lines should be glued. $R^{e,\nu}$ is transverse to φ_{geod} and the induced coorientation is shown in red. The induced orientation of the horizontal boundary of $R^{e,\nu}$ (red) is opposed to the orientation of the flow (black). Thus the surfaces we will construct are negative Birkhoff cross sections.

boundary in a vertex of γ is empty if only if two adjacent edges are cooriented in a direction, and two others in the opposite direction: this means that ν is Eulerian around v. Conversely, if ν is Eulerian, then up to rotation there are two local configurations around v (that we called alternating and transparent), and one checks that in both cases, the vertical boundary is empty (see the left parts of Figures 5 and 6). □

When ν is Eulerian, the complex $S^\times(\nu)$ is not a topological surface if ν has some transparent points: as depicted on Figure 6, there are edges adjacent to four faces. But it is the only obstruction and we can desingularize such segments. Also if we want a smooth surface, we have to smooth $S^\times(\nu)$ is a neighborhood of the fibers of the double points. In this way, we obtain a smooth surface, transverse to φ_{geod}.

Definition 24. For ν an Eulerian coorientation, the associated **BB-surface** is the surface $S^{BB}(\nu)$ obtained from $S^\times(\nu)$ by desingularizing and smoothing the fibers of the double points of γ (see the right parts of Figures 5 and 6).

For example, the BB-surface associated to a Birkhoff coorientation (Example 14) is isotopic to the construction suggested by Birkhoff [Bir17] and popularized by Fried [Fri83]. Also the BB-surface associated to a Brunella coorientation (Example 15) has been introduced by Brunella [Bru94, Description 2].

3.b. **Asymptotic cycles and existence of sections.** The question whether a given vector field admits a global section (*i.e.*, with empty boundary) has been given a very satisfactory answer by Schwarzmann and Fuller [Sch57, Ful65], then expanded by Fried [Fri82].



FIGURE 5. On the left, the complex $S^\times(v)$ around the fiber of an alternating double point of γ. Every point of the fiber of v is adjacent to exactly two rectangles. On the right the surface $S^{BB}(v)$ is obtained by smoothing $S^\times(v)$.

A preliminary remark: if two surfaces S_1 and S_2 in a manifold M are global sections to a flow ϕ and they are homologous, then they are isotopic, and the isotopy is realized by the flow. Indeed[4] one can consider the infinite cyclic covering of $\hat{M} \to M$ associated to the morphism $\pi_1(M) \to \mathbb{Z}$ given by the intersection with $[S_1] = [S_2]$. Then S_1 and S_2 lift into $\mathbb{Z}$ disjoint copies $t^n\hat{S}_1$ and $t^n\hat{S}_2$ in $\hat{M}$, all transverse to the lift of the flow. Now following the flow starting from $\hat{S}_1$, one reaches $\hat{S}_2$, so we have a surjective map $\hat{S}_1 \to \hat{S}_2$ of local degree 1, and since $\hat{S}_1$ is transverse to the flow it is of total degree 1. Similarly we have a surjection $\hat{S}_2 \to \hat{S}_1$ of local degree 1. By composing the two, we get of surjection $\hat{S}_1 \to \hat{S}_1$ of total degree 1, hence a bijection. Therefore the maps $\hat{S}_1 \to \hat{S}_2$ and $\hat{S}_2 \to \hat{S}_1$ are actually bijections, and the flow hence induces an isotopy $\hat{S}_1 \to \hat{S}_2$. Projecting back in M, we obtained the desired isotopy $S_1 \to S_2$.

For X a vector field in a compact manifold M, we denote by $k_X(p,t)$ a closed curve obtained by concatenating the piece of orbit $\phi^{[0,t]}(p)$ starting at p of length t with an arc connecting $\phi^t(p)$ to $\phi^0(p)$ of bounded length. The class $[k_X(p,t)]$ in $\mathrm{H}_1(M;\mathbb{Z})$ then depends on the choice of the closing segment, but only in a bounded way, so that the limit $\lim_{t\to\infty} \frac{1}{t}[k_X(p,t)]$, if it exists, does not depend on this choice. An **asymptotic cycle** of X is then the limit of a sequence of the form $\{\frac{1}{t_n}[k_X(p_n,t_n)] \,|\, p_n \in M, t_n \to \infty\}$ in $\mathrm{H}_1(M;\mathbb{R})$. The set of asymptotic cycles is denoted $\mathcal{S}_X$.

[4]This mimics the folklore argument in knot theory that the fiber of a fibration minimizes the genus, but it is not so easy to find a reference of this statement.



FIGURE 6. On the left, the complex $S^{\times}(v)$ around the fiber of a transparent double point of γ. Every point of the fiber of v is adjacent to an even number of rectangles. On the right the surface $S^{BB}(v)$ is obtained by desingularizing $S^{\times}(v)$ on the portion of the fiber where four rectangles meet.

Sullivan [Sul76] reinterpreted it by showing that every X-invariant measure μ induces a *foliated cycle* c_μ that is actually a positive combination of asymptotic cycles.

Theorem 25. [Sch57, Ful65] *A vector field X on a closed M admits a global section whose homology class is $\sigma \in \mathrm{H}_2(M, \partial M; \mathbb{Z})$ if and only σ intersects positively every asymptotic cycle, namely for every $c \in \mathcal{S}_X$ one has $\langle \sigma, c \rangle > 0$.*

This theorem is beautiful, but unfortunately, for many vector fields X, the point 0 belongs to $\mathrm{Conv}(\mathcal{S}_X)$, so that X admits no global section at all. This is where Birkhoff cross sections come in.

3.c. **Classes of surfaces with given boundary.** Now we work in our restricted setting: Σ is a negatively curved surface, γ is a finite collection of periodic geodesics and $\overleftrightarrow{\gamma}$ denotes the antithetic lift of γ. In order to apply Theorem 25 for finding Birkhoff cross sections, we need to work in the complement $\mathrm{T}^1\Sigma \setminus \overleftrightarrow{\gamma}$ and in particular to determine the space $\mathrm{H}_2(\mathrm{T}^1\Sigma, \overleftrightarrow{\gamma}; \mathbb{Z})$. In this section we show that the homology classes of surfaces bounded by $-\overleftrightarrow{\gamma}$ form an affine space and we give a canonical origin to this space.

Lemma 26. *The sequence $0 \to \mathrm{H}_2(\mathrm{T}^1\Sigma; \mathbb{Z}) \xrightarrow{i} \mathrm{H}_2(\mathrm{T}^1\Sigma, \overleftrightarrow{\gamma}; \mathbb{Z}) \xrightarrow{\partial} \mathrm{H}_1(\overleftrightarrow{\gamma}; \mathbb{Z})$, where the first map is the inclusion map and the second is the boundary map, is exact.*[5]

[5] An erroneous version of this statement is in [Fri82, Lemma 6], where it is claimed that the boundary map is surjective and admits a section. It is not true in general, unless the manifold is a homology sphere.

Proof. This is just a part of the long exact sequence associated to the pair $(\mathrm{T}^1\Sigma, \overleftrightarrow{\gamma})$, see [Hat02, Thm 2.16], where we note that $\mathrm{H}_2(\overleftrightarrow{\gamma};\mathbb{Z})$ is zero. □

The homology classes of those surfaces whose boundary is $-\overleftrightarrow{\gamma}$ correspond to the preimages by ∂ of the point $\{-1,-1,\dots,-1\} \in \mathrm{H}_1(\overleftrightarrow{\gamma};\mathbb{Z}) \simeq \mathbb{Z}^{2|\gamma|}$. Hence they form an affine space directed by $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{Z})$. Indeed given two surfaces with the same boundary, their difference induces a class in $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{Z})$. Now using the fact that $\mathrm{T}^1\Sigma$ is a circle bundle with non-zero Euler class, we get $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{Z}) \simeq \mathrm{H}_1(\Sigma)$: a non-trivial class in $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{Z})$ can be represented by the set of the fibers over a cycle in $\mathrm{H}_1(\Sigma)$.

From the previous discussion we deduce that if we are given a explicit surface S_0 bounded by $-\overleftrightarrow{\gamma}$, the classes of the other surfaces bounded by $-\overleftrightarrow{\gamma}$ differ from $[S_0]$ by a class in $\mathrm{H}_1(\Sigma)$. In our context, there is a natural choice of such an origin, for which the computation of the intersection numbers with asymptotic cycles of the geodesic flow will be easy. We denote by $S^\times_\pm$ the rational chain in $\mathrm{H}_2(\mathrm{T}^1\Sigma \setminus \overleftrightarrow{\gamma}, \overleftrightarrow{\gamma};\mathbb{Q})$ that is half the sum of all rectangles of the form $R^{e,\nu}$ (see Figure 7):

$$S^\times_\pm := \frac{1}{2} \sum_{e\in\gamma, \nu_e = \pm} R^{e,\nu_e}.$$

In other words, we consider the set of all tangent vectors base at points of γ. Remember that every rectangle is cooriented by the geodesic flow, hence oriented. Therefore, $S^\times_\pm$ is also oriented. Its boundary is then exactly $-\overleftrightarrow{\gamma}$. The chain $S^\times_\pm$ is not a surface since the fibers of the double points of γ are singular. Its double is an integer class. As it is rational the class $\sigma_\pm := [S^\times_\pm]$ might not be realized by a surface.[6]

Lemma 27. *For $\underrightarrow{\alpha}$ an oriented periodic geodesic on Σ that is not a component of γ, the algebraic intersection $\langle \sigma_\pm, \vec{\alpha}\rangle$ is equal to $+\frac{1}{2}|\{\underrightarrow{\alpha} \cap \gamma\}|$.*

This lemma appears in a different form in [DIT15] where it is used to prove that the linking number of two collections $\overleftrightarrow{\gamma_1}, \overleftrightarrow{\gamma_2}$ in $\mathrm{T}^1\Sigma$ is actually equal to $|\{\gamma_1 \cap \gamma_2\}|$.

Proof. Since $S^\times_\pm$ is positively transverse to the geodesic flow, all intersection points of $\vec{\alpha}$ with $S^\times_\pm$ counts positively. Since every rectangle has coefficients $\frac{1}{2}$ in $S^\times_\pm$, every intersection point contributes for $+\frac{1}{2}$ to the algebraic intersection. Finally $\vec{\alpha}$ intersects $S^\times_\pm$ exactly in the fiber of the intersection points of $\underrightarrow{\alpha}$ and γ. □

3.d. **Proofs of Theorems C and D.** Denote by $\mathcal{S}_{\overleftrightarrow{\gamma}} \subset \mathrm{H}_1(\mathrm{T}^1\Sigma \setminus \overleftrightarrow{\gamma};\mathbb{R})$ the set of asymptotic cycles of the geodesic flow φ_{geod} restricted to $\mathrm{T}^1\Sigma \setminus \overleftrightarrow{\gamma}$. Also denote by π the canonical projection from $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{R})$ to $\mathrm{H}_1(\Sigma;\mathbb{R})$. The next statement is the key to our main result.

Lemma 28. *A class $\sigma \in \mathrm{H}_2(\mathrm{T}^1\Sigma, \overleftrightarrow{\gamma};\mathbb{R})$ intersects positively every element of $\mathcal{S}_{\overleftrightarrow{\gamma}}$ if and only if the class $\pi(\sigma - \sigma_\pm) \in \mathrm{H}_1(\Sigma;\mathbb{R})$ lies in the interior of $\frac{1}{2}B^*_{\|\cdot\|_\gamma}$.*

[6] Actually, $\sigma_\pm$ is realized by a surface if and only if $[\gamma]_2$, the class of γ with $\mathbb{Z}/2\mathbb{Z}$-coefficients, is 0. In this case, the homology class of Birkhoff's coorientation ν_B (Example 14) is 0, and $S^{BB}(\nu_B)$ lies in the class $\sigma_\pm$. Also the class $\sigma_\pm$ is equal to $\frac{1}{2}[S^{BB}(\nu) + S^{BB}(-\nu)]$ for every Eulerian ν. Hence it is always realized as the average of two surfaces without any assumption on $[\gamma]_2$.



FIGURE 7. The 2-chain $S^{\times}_{\pm}$ is half of the sum of all rectangles R^{e,ν_e}. It is cooriented by the geodesic flow, hence oriented (in red). Its boundary, taking orientations into account, is then $-\overleftrightarrow{\gamma}$.

Proof. By the shadowing property for pseudo-Anosov flows, the projectivization of $\mathcal{S}_{\overleftrightarrow{\gamma}}$ is the convex hull of the cycles given by periodic orbits. Hence it is enough to estimate the intersection of σ with all periodic orbits of φ_{geod}.

We use the bracket to denote the intersection, and the index reminds the space where the objects live. For every periodic orbit $\vec{\alpha}$ of φ_{geod}, by Lemma 27, we have

$$\begin{aligned}\langle \sigma, \vec{\alpha}\rangle_{\mathrm{T}^1\Sigma\setminus\overleftrightarrow{\gamma}} &= \langle \sigma - \sigma_\pm, \vec{\alpha}\rangle_{\mathrm{T}^1\Sigma\setminus\overleftrightarrow{\gamma}} + \langle \sigma_\pm, \vec{\alpha}\rangle_{\mathrm{T}^1\Sigma\setminus\overleftrightarrow{\gamma}} \\ &= \langle \sigma - \sigma_\pm, \vec{\alpha}\rangle_{\mathrm{T}^1\Sigma\setminus\overleftrightarrow{\gamma}} + \frac{1}{2}|\{\underrightarrow{\alpha} \cap \gamma\}| \\ &= \langle \pi(\sigma - \sigma_\pm), \underrightarrow{\alpha}\rangle_\Sigma + \frac{1}{2}|\{\underrightarrow{\alpha} \cap \gamma\}|.\end{aligned}$$

Hence $\langle \sigma, \vec{\alpha}\rangle_{\mathrm{T}^1\Sigma\setminus\overleftrightarrow{\gamma}}$ is positive if and only if $-\langle \pi(\sigma - [S^{\times}_{\pm}]), \underrightarrow{\alpha}\rangle_\Sigma$ is smaller than $\frac{1}{2}|\{\underrightarrow{\alpha} \cap \gamma\}|$.

Now the term $-\langle \pi(\sigma - \sigma_\pm), \underrightarrow{\alpha}\rangle_\Sigma$ depends only on the class $[\underrightarrow{\alpha}] \in \mathrm{H}_1(\Sigma;\mathbb{Z})$, while the term $\frac{1}{2}|\{\underrightarrow{\alpha} \cap \gamma\}|$ is larger that $\frac{1}{2}\|([\underrightarrow{\alpha}])\|_\gamma$, with equality if $\underrightarrow{\alpha}$ is $\|\cdot\|_\gamma$-minimizing. Hence the inequality $-\langle \pi(\sigma - \sigma_\pm), \underrightarrow{\alpha}\rangle_\Sigma \leq \frac{1}{2}|\{\underrightarrow{\alpha} \cap \gamma\}|$ is equivalent to $-\langle \pi(\sigma - \sigma_\pm), [\underrightarrow{\alpha}]\rangle_\Sigma \leq \frac{1}{2}\|([\underrightarrow{\alpha}])\|_\gamma$.

Therefore σ intersects positively every element of $\mathcal{S}_{\overleftrightarrow{\gamma}}$ if and only if for every class $a \in \mathrm{H}_1(\Sigma;\mathbb{Z})$ we have $-\langle \pi(\sigma - \sigma_\pm), a\rangle_\Sigma \leq \frac{1}{2}\|a\|_\gamma$, which means exactly that the point $-\pi(\sigma - \sigma_\pm)$ belongs to $\frac{1}{2}B^*_{\|\cdot\|_\gamma}$. Since the latter is symmetric about the origin, this amounts to $\pi(\sigma - \sigma_\pm)$ belonging to $\frac{1}{2}B^*_{\|\cdot\|_\gamma}$. □

As a byproduct of the proof, we obtain that a class $\sigma \in \mathrm{H}_2(\mathrm{T}^1\Sigma, \overleftrightarrow{\gamma};\mathbb{R})$ intersects non-negatively every asymptotic cycle if and only if $\pi(\sigma - \sigma_\pm) \in \mathrm{H}_1(\Sigma;\mathbb{R})$ lies in the closed unit ball $\frac{1}{2}B^*_{\|\cdot\|_\gamma}$.

Proof of Theorem C. For ν an Eulerian coorientation, we consider the surface $S^{BB}(\nu)$ (Definition 24). By construction it is transverse to the geodesic flow. One easily checks that every rectangle of the form $R^{e,\nu}$ contributes to -1 to the Euler characteristics, hence $\chi(S^{BB}(\nu))$ is $-|E(\gamma)|$. Since γ is a graph of degree 4, one has $|E(\gamma)| = 2|V(\gamma)|$, so that $\chi(S^{BB}(\nu)) = -2|V(\gamma)|$.

Now if ν_1 and ν_2 are cohomologous, the class $[S^{BB}(\nu_1) - S^{BB}(\nu_2)] \in \mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{Z})$ projects by π onto $[\nu_1 - \nu_2] = 0$. Since π is actually an isomorphism, $[S^{BB}(\nu_1) - S^{BB}(\nu_2)] = 0$, which in turn implies $[S^{BB}(\nu_1)] = [S^{BB}(\nu_2)]$ in $\mathrm{H}_2(\mathrm{T}^1\Sigma, \overleftrightarrow{\gamma};\mathbb{Z})$. Now since $S^{BB}(\nu_1)$ and $S^{BB}(\nu_2)$ are both transverse to φ_{geod} and homologous, the flow actually realizes an isotopy between them. □

Proof of Theorem D. Let us first treat the case when Σ has genus 2 or higher. Lemma 26 and the paragraph after implies that real homology classes of surfaces bounded by $-\overleftrightarrow{\gamma}$ form an affine space directed by $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{R})$. The class $\sigma_\pm$ also defined in 3.c gives a canonical origin to this space. It is a half-integer class, and its double $2\sigma_\pm$ is congruent to $[\gamma]_2$ mod 2. Therefore the double of all integer classes correspond to the sublattice of $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{Z})$ of those points congruent to $[\gamma]_2$ mod 2.

Now we have to determine which of these integer classes yield Birkhoff cross sections. By Schwarzmann-Fuller Theorem 25, a class σ contains a Birkhoff cross section if and only if it intersects positively every asymptotic cycles. By Lemma 28 this means that the difference $\sigma - \sigma_\pm$ lies inside $\frac{1}{2}B^*_{\|\cdot\|_\gamma}$, or equivalently that $2(\sigma - \sigma_\pm)$ lies inside $B^*_{\|\cdot\|_\gamma}$.

Now surfaces that are transverse to φ_{geod} correspond to homology classes that intersects non negatively every asymptotic cycle, allowing certain intersection to be zero. This means that the boundary of $B^*_{\|\cdot\|_\gamma}$ is now authorized. This end the proof for surface of genus at least 2.

For the case of the torus, the only difference is that the bundle $\mathrm{T}^1\Sigma$ is trivial, *i.e.,* of the form $\Sigma \times \mathbb{S}^1$. Therefore we no longer have $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{R}) \simeq \mathrm{H}_1(\Sigma;\mathbb{R})$, but instead $\mathrm{H}_2(\mathrm{T}^1\Sigma;\mathbb{R}) \simeq \mathrm{H}_1(\Sigma;\mathbb{R}) \times \mathbb{R}$, since the fibers are no longer boundaries. However this extra $\mathbb{R}$-factor does not change the proof, since all asymptotic cycles of the geodesic flow on the flat torus are actually horizontal, meaning that the extra coordinate is zero. Hence the positivity condition depends only on the coordinate in $\mathrm{H}_1(\Sigma;\mathbb{R})$, and all the arguments can be translated. □

4. Questions

On intersection norms. If Σ is a flat torus, then the minimal intersection is always realized by geodesics, which are unique in their homology class. Hence if γ is the union of k geodesics $\gamma_1, \dots, \gamma_k$, then $i_\gamma(\alpha) = \sum_{i=1}^k i_{\gamma_i}(\alpha)$. This implies that the dual ball B^*_γ coincides with the Minkowski sum $B^*_{\gamma_1} + \cdots + B^*_{\gamma_k}$. Since the segment $[-1,1] \times \{0\} \subset \mathbb{R}^2$ is the dual unit ball $B^*_{\|\cdot\|_\gamma}$ for γ the vertical circle on the torus, every segment containing 0 in the middle is the dual unit ball of some closed

circle on the torus. Therefore every convex polygon in $\mathbb{R}^2$ whose vertices are integral and congruent mod 2 is of the form $B^*_{\|\cdot\|_\gamma}$ for some γ. In higher dimension the situation is probably more intricate.

Question 29. Which polyhedra of $\mathbb{R}^{2g}$ with integer vertices can be realized as the dual unit ball $B^*_{\|\cdot\|_\gamma}$ for some γ in Σ_g?

Also, if Σ is a torus and γ is a union of geodesics, then the above remarks imply that the number of self-intersection points of γ is exactly $1/4$ of the area of $B^*_{\|\cdot\|_\gamma}$ (check on Figure 1). Is there an analog statement in higher genus?

Question 30. Which information concerning γ can be read on $B^*_{\|\cdot\|_\gamma}$? Is the number of self-intersection points of γ a certain function defined on $B^*_{\|\cdot\|_\gamma}$?

This information is interesting since the this number is exactly the opposite of the Euler characteristic of every Birkhoff cross section bounded by $\overleftrightarrow{\gamma}$. Note the the number of self-intersection points is homogenous of degree 2, so we should look for degree 2 functions on polyhedra in $\mathbb{R}^{2g}$: does it correspond to some symplectic capacity?

Motivated by our application we only defined the intersection norm for a collection of immersed curves, but one can directly extend it for an arbitrary embedded graph. One can wonder which properties extend to this case and which information on the embedded graphs are encoding in this norm. For example when the graph is Eulerian (*i.e.,* all vertices have even degree) the connection with Eulerian coorientations remains.

On Birkhoff cross sections. Our constructions and our classification result deal only with Birkhoff cross sections bounded by an **amphithetic** collection of periodic orbits of the geodesic flow, that is, invariant under the involution $(p, v) \mapsto (p, -v)$. However the only restriction *a priori* for being the boundary of a Birkhoff cross section is to be a boundary, that is, to be null-homologous. Our results here say nothing about the classification, or even the existence, of Birkhoff cross sections with arbitrary null-homologous boundary. In this case, the theory of Schwarzmann-Fuller-Thurston-Fried and the remarks of Sections 3.b and 3.c still apply, so that these sections still correspond to the point inside a certain polytope in $\mathrm{H}^1(\Sigma; \mathbb{R})$. However we have no analog for the coorientations and the explicit constructions derived from them.

Question 31. Is there a natural generalization of the polytope $B^*_{\|\cdot\|_\gamma}$ to non-amphithetic collections $\vec{\gamma}$ of orbits of the geodesic flow φ_{geod}, so that integer points in this polytope classify surfaces bounded by $\vec{\gamma}$ and transverse to φ_{geod}?

In the case of the flat torus, this question is answered in [Deh15a, Thm 3.12] where a polygon $P_{\vec{\gamma}}$ classifying transverse surfaces bounded by $\vec{\gamma}$ is defined for *every* null-homologous collection $\vec{\gamma}$.

What would probably unlock the situation in the higher genus case would be to have, for every null-homologous collection $\vec{\gamma}$, *one* explicit surface bounded by $\vec{\gamma}$ (not necessarily transverse), that is, a analog of $\sigma_\pm$ when $\vec{\gamma}$ is not antithetic. Such an explicit point allows to compute its intersection with every other periodic orbit $\vec{\alpha}$ of φ_{geod}. These intersection numbers are all we need in order to describe explicitly the asymptotic directions of φ_{geod} in $\mathrm{T}^1\Sigma \setminus \vec{\gamma}$. Generalising the constructions of [Deh15b] is a possibility here.

More generally, one can wonder whether there exists a generalization to all flows of the intersection norm $\|\cdot\|_\gamma$ in the following sense:

Question 32. For every 3-dimensional flow X, is there an object that describes all isotopy classes of Birkhoff cross sections?

A starting point would be to try with an Anosov flow that is not the geodesic flow, and see whether Gauss linking forms [Ghy09] could play this role.

References

[Ano67] Anosov Dmitri V.: Geodesic flows on closed Riemannian manifolds with negative curvature, *Proc. Steklov Inst. Math.* **90** (1967), 235pp.

[Bir17] Birkhoff George D.: Dynamical systems with two degrees of freedom, *Trans. Amer. Math. Soc.* **18** (1917), 199–300.

[Bru94] Brunella Marco: On the discrete Godbillon-Vey invariant and Dehn surgery on geodesic flow, *Ann. Fac. Sc. Toulouse Sér. 6* **3** (1994), 335–344.

[Cal06] Calegari Danny: Universal circles for quasigeodesic flows, *Geom. Topol.* **10** (2006), 2271–2298.

[Deh15a] Dehornoy Pierre: Geodesic flows, left-handedness, and templates, *Algebr. Geom. Topol.* **15** (2015), 1525–1597.

[Deh15b] Dehornoy Pierre: Genus one Birkhoff cross sections for geodesic flows, *Ergod. Theory Dynam. Systems* **35** (2015), 1795–1813.

[Fri79] Fried David: Fibrations over $\mathbb{S}^1$ with pseudo-Anosov monodromy, Exposé 14, in *Fathi, Laudenbach, Poenaru*: *Travaux de Thurston sur les surfaces*, *Astérisque* **66-67** (1979).

[Fri82] Fried David: The geometry of cross sections to flows, *Topology* **21** (1982), 353–371.

[Fri83] Fried David: Transitive Anosov flows and pseudo-Anosov maps, *Topology* **22** (1983), 299–303.

[Ful65] Fuller Francis Brock: On the surface of section and periodic trajectories, *Am. J. Math.* **87** (1965), 473–480.

[Ghy09] Ghys Étienne: Right-handed vector fields and the Lorenz attractor, *Japan. J. Math.* **4** (2009), 47–61.

[Gro76] Gromov Mikhaïl: Three remarks on geodesic dynamics and fundamental group, *preprint SUNY* (1976), reprinted in *Enseign. Math.* **46** (2000), 391–402.

[Had1898] Hadamard Jacques: Les surfaces à courbures opposées et leurs lignes géodésiques, *J. Math. Pures Appl.* **4** (1898), 27–74.

[Hat02] Hatcher Allen: *Algebraic topology*, Cambridge University Press (2002), 550+xii pp.

[Hof60] Hoffman Alan J.: Some recent applications of the theory of linear inequalities to extremal combinatorial analysis, in: *Combinatorial Analysis* (R. Bellman, M. Hall Jr, eds.) [*Proceedings of Symposia in Applied Mathematics*, Volume X] (1960), 113–127.

[Mos92] Mosher Lee: Dynamical systems and the homology norm of a 3-manifold II, *Invent. Math.* **107** (1992), 243–281.

[Sal16] de la Salle Mikael: Another proof of Thurston's theorem for norms taking integer values on the integer lattice, arXiv:1604.01578

[Sch03] Schrijver Alexander: *Combinatorial Optimization, Polyhedra and Efficiency*, Springer (2003).

[Sch57] Schwarzmann Sol: Asymptotic cycles, *Ann. Math.* **66** (1957), 270–284.

[Sul76] Sullivan Dennis: Cycles for the dynamical study of foliated manifolds and complex manifolds, *Invent. Math.* **36** (1976), 225–255.

[Thu86] Thurston William: A norm for the homology of three-manifolds, *Mem. Amer. Math. Soc.* **339** (1986), 99–130.

[DIT15] Duke William, Imamöglu Özlem, and Tóth Árpád: Linking numbers and modular cocycles, *preprint*.

[Whi34] Whitney Hassler: Analytic extensions of Differentiable functions defined in closed sets, *Trans. Amer. Math. Soc.* **36** (1934), 63–89.

[Yaz16] Yazdi Mehdi: On Thurston's Euler class one conjecture, arXiv:1603.03822

IMPA, Estrada Dona Castorina 110, 22460-320 Rio de Janeiro, Brasil
marcossa@impa.br

Univ. Grenoble Alpes, IF, F-38000 Grenoble, France
CNRS, IF, F-38000 Grenoble, France

pierre.dehornoy@ujf-grenoble.fr
http://www-fourier.ujf-grenoble.fr/~dehornop/